

Adherence ^[1]

Submitted on May 12, 2014



Table of Contents

- [Strong Drugs to Fight HIV](#)
- [The Importance of Adherence](#)
- [Starting and Choosing HIV Drugs](#)
- [Barriers to Adherence for Women](#)
- [Overcoming Barriers](#)
- [Adherence Tips](#)
- [Finding Support](#)

Strong Drugs to Fight HIV

The many advances in HIV treatment in the past 20 years have led to the approval of over 25 drugs that fight HIV, as well as seven combination pills that contain two or more HIV drugs. This is a remarkable accomplishment. Most importantly, using these drugs in effective combinations is helping many people control their HIV and live longer and healthier lives.

In order for HIV drugs to do their job properly, you need to do yours! It is very important that you take your drugs exactly as they are prescribed. That means taking the correct amount (dose) at the right times, and as directed (e.g., with any food or medication restrictions). This is called adherence.

Despite the improvements in HIV drug treatment (such as fewer [side effects](#) [2], fewer restrictions, and easier dosing), taking HIV drugs is still a daily, lifelong commitment. While this can make adherence challenging for many people living with HIV (HIV+), the benefits are well worth the effort.

The Importance of Adherence

When you take a drug, it gets processed by your body and enters your blood stream. The HIV drugs need to stay in your blood at certain levels to fight HIV. If the level falls too low, there is not enough drug to do the job.

When you take your pills on schedule, you keep the right level of the drug in your body. If you do not take your HIV drugs on schedule, drug levels in your blood will drop and HIV will be able to make copies of itself (multiply). When HIV multiplies, your [viral load](#) [3] increases and your [CD4 cell](#) [4] count goes down.

When HIV makes copies of itself, it sometimes makes changes, called mutations. These mutations can help the virus survive, even when you are taking your HIV medication. This is called [resistance](#) [5]. When HIV becomes resistant to a drug you are taking, that drug will stop working. At that point, you will probably have to switch HIV drugs.

Resistance to one drug can sometimes cause resistance to other drugs you have not taken. This is called cross-resistance. Resistance often develops to other drugs in the same class. If you were taking one non-nucleoside reverse transcriptase inhibitor (NNRTI) ? say Sustiva (efavirenz) ? and you developed resistance to Sustiva, you would likely develop resistance to another NNRTI, like Intelence (etravirine). Resistance can affect your treatment choices in the future because fewer drugs will work well against your virus.

The best way to prevent resistance is to stick closely to your medication schedule. Some studies suggest that taking your HIV drugs exactly as prescribed about 95 percent of the time is needed to get the most benefit from HIV treatment. That means missing very few doses.

Starting and Choosing HIV Drugs

There are a number of things to think about when deciding whether to start HIV drug treatment. Two of those things are your CD4 count and viral load.

It is also important to think about whether you feel able to commit to taking your drugs correctly. If you do not feel ready, talk it over with someone who knows about HIV. Your health care provider or the staff at an AIDS service organization (ASO) can give you accurate information to answer your questions and address your concerns.

Barriers to Adherence for Women

Many women find it difficult to be adherent to their HIV medication for the following reasons:

- A busy lifestyle
- Not having disclosed [6] their HIV status to work colleagues, friends, lovers, or family
- Depression [7], a leading cause of non-adherence, is more common in HIV+ women than HIV+ men
- Other life stressors, such as childcare or parenting issues
- Current or past problems with side effects
- Active substance use [8] or problems with alcohol
- In the US, women are less likely to have health insurance than men. Without health insurance, women are unlikely to be able to access adequate care and HIV drugs
- Many HIV+ women do not have safe or stable housing; this presents additional barriers to getting treatment [9]
- Past or current experience of violence or trauma. Women who have been abused are four times more likely not to adhere to their HIV drugs. Globally, violence against women is very common. For more information, see our article on Violence Against Women and HIV [10].

Overcoming Barriers

Despite the many demands in women?s lives, numerous studies have found that women are just as likely to adhere to HIV regimens as men, especially when women have longstanding and trusting relationships with their health care providers.

Even though it may be embarrassing, it is important to tell your health care provider about the

number of times you have missed doses or did not take your HIV drugs correctly. He or she may suggest a change in your dosing schedule or drug regimen that makes it easier. Newer HIV drugs require fewer pills per day and have fewer food restrictions. There are also new ways to combine older drugs that make them easier to take.

Adherence Tips

- Understand that the medications will help you fight the virus and stay well. If you do not think they will, you will not bother taking your pills correctly. If you have any doubts, speak to your health care provider or staff at an AIDS service organization (ASO).
- Some women find that connecting their pill-taking with a higher purpose or good intention helps. This intention could be spiritual in nature, or something like staying healthy for the sake of family, friends, or others served by your unique gifts.
- Use a daily activity, one that you do every day without fail (like waking up in the morning, walking your dog, or going to bed at night), to remind you to take your pills. When it is time to do that activity, you will know that it is also time to take your pills. If you have children who take medicines or vitamins, you may consider taking your HIV drugs when they take their medicine so the whole family stays healthy together.
- If you do not want others to see you taking your pills, quietly slip away to a private area or the bathroom. If that will not work, say the medications are for another health problem or that they are vitamins.
- If you think substance use or mental health issues are preventing you from taking your medications correctly, talk to your health care provider or case manager so they can get you help. There are good treatments available.
- Take advantage of tools available from your clinic or pharmacy such as pillboxes, calendars, diaries, and beepers to help you remember to take your medications on time.
- Plan ahead for refills or trips so you do not run out of any medications.

Finding Support

Adherence is hard work and takes a lot of commitment. It helps to have other people on your side. One way to do this is to put together a support network. Your health care provider is one of the most important people in your network. Talk openly with him or her about how to fit HIV treatment into your lifestyle.

There are many other sources of information and support available to women who are taking or thinking about taking HIV drugs. If you can, include family, friends, case managers, treatment educators, and counselors in your network. You can also get involved with your local ASO or a support group. These are places where you will be able to ask questions, get good tips, and share experiences with others who understand what it is like to live with HIV. When you are feeling discouraged, turn to your network for support and encouragement.

Also, try to remember the big picture. It can be hard to take pills every day, but easier when you know why you are doing it. You want to be adherent to give your HIV drugs the best chance of working to keep you healthy. This will allow you to focus on the things you care about ? including the important people in your life like your children, partners, family, and friends.

Tags:

- [women and HIV adherence treatment](#) [11]
- [schedule adherence](#) [12]
- [HIV adherence treatment](#) [13]
- [medication adherence](#) [14]
- [adherence drug](#) [15]
- [adherence](#) [16]
- [HIV adherence](#) [17]
- [HIV resistance](#) [18]
- [HIV drugs](#) [19]
- [staying adherent](#) [20]
- [adherence tips](#) [21]
- [women adherence](#) [22]
- [HIV women adherence](#) [23]
- [taking HIV drugs](#) [24]
- [skipping HIV drugs](#) [25]
- [reminder to take HIV meds](#) [26]

Additional Resources

Select the links below for additional material related to adherence.

[Staying on Schedule: Tips for Taking your HIV Medicines \(NY State\)](#) [27]

[Adherence to Treatment for HIV: Training Curricula in India \(EngenderHealth\)](#) [28]

[The Importance of Adhering to your Treatment Regimen \(AIDSmeds\)](#) [29]

[What is Treatment Adherence? \(AIDSinfo\)](#) [30]

[Following an HIV Treatment Regimen \(The Body/AIDSinfo\)](#) [31]

[Adherence \(AIDS InfoNet\)](#) [32]

[Six Reasons Why People Skip Their HIV Meds \(The Body\)](#) [33]

[Treatment Adherence \(AIDS.gov\)](#) [34]

[Financial stress?impact on HIV adherence, HCV, and prescribing patterns \(CATIE\)](#) [35]

[Making Your Treatment Work Long-Term \(CATIE\)](#) [36]

- [Sign Up / Login](#)
- [My Account](#)
- [HIV Information](#)
- [A Girl Like Me](#)
- [Partners](#)
- [Who We Are](#)
- [Terms](#)
- [Privacy](#)
- [Contact](#)



@2015 thewellproject. All rights reserved.

Source URL: <http://www.thewellproject.org/hiv-information/adherence-0>

Links:

- [1] <http://www.thewellproject.org/hiv-information/adherence-0>
- [2] <http://www.thewellproject.org/hiv-information/side-effects>
- [3] <http://www.thewellproject.org/hiv-information/women-and-viral-load>
- [4] <http://www.thewellproject.org/hiv-information/understanding-cd4-cells-and-cd4-cell-tests>
- [5] <http://www.thewellproject.org/hiv-information/resistance>
- [6] <http://www.thewellproject.org/hiv-information/disclosure-and-hiv>
- [7] <http://www.thewellproject.org/hiv-information/depression-women-and-hiv>
- [8] <http://www.thewellproject.org/hiv-information/substance-abuse-and-addiction>
- [9] <http://www.thewellproject.org/hiv-information/getting-hiv-drugs-us>
- [10] <http://www.thewellproject.org/hiv-information/violence-against-women-and-hiv>
- [11] <http://www.thewellproject.org/tags/women-and-hiv-adherence-treatment>
- [12] <http://www.thewellproject.org/tags/schedule-adherence>
- [13] <http://www.thewellproject.org/tags/hiv-adherence-treatment>
- [14] <http://www.thewellproject.org/tags/medication-adherence>
- [15] <http://www.thewellproject.org/tags/adherence-drug>
- [16] <http://www.thewellproject.org/tags/adherence>
- [17] <http://www.thewellproject.org/tags/hiv-adherence>
- [18] <http://www.thewellproject.org/tags/hiv-resistance>
- [19] <http://www.thewellproject.org/tags/hiv-drugs>
- [20] <http://www.thewellproject.org/tags/staying-adherent>
- [21] <http://www.thewellproject.org/tags/adherence-tips>
- [22] <http://www.thewellproject.org/tags/women-adherence>
- [23] <http://www.thewellproject.org/tags/hiv-women-adherence>
- [24] <http://www.thewellproject.org/tags/taking-hiv-drugs>
- [25] <http://www.thewellproject.org/tags/skipping-hiv-drugs>
- [26] <http://www.thewellproject.org/tags/reminder-take-hiv-meds>
- [27] <http://www.health.ny.gov/publications/9236.pdf>
- [28] <http://www.engenderhealth.org/our-countries/asia-near-east/india-publications.php>
- [29] http://www.aidsmeds.com/articles/Adherence_7420.shtml
- [30] http://aidsinfo.nih.gov/contentfiles/WhatIsTreatmentAdherence_FS_en.pdf
- [31] <http://www.thebody.com/content/50168/following-an-hiv-treatment-regimen.html>
- [32] http://www.aidsinfonet.org/fact_sheets/view/405
- [33] <http://www.thebody.com/content/63738/reasons-why-people-skip-their-meds.html?getPage=1>
- [34] <http://aids.gov/hiv-aids-basics/just-diagnosed-with-hiv-aids/treatment-options/medication-adherence/>
- [35] <http://www.catie.ca/en/catienews/2012-10-30/financial-stress-impact-hiv-adherence-hcv-and-prescribing-patterns>
- [36] <http://www.catie.ca/en/practical-guides/hiv-drug-treatment/4-once-youve-started/4-2>