

Considerations Before Starting HIV Treatment

Last updated: March 30, 2021

Together, we can change the course of the HIV epidemic...one woman at a time.

#onewomanatatime

#thewellproject

www.thewellproject.org



- If you're living with HIV, treatment with HIV drugs can improve your quality of life and help you stay healthier longer
- Starting treatment is a big decision





Preparing for HIV Treatment

- To get the most benefit from HIV drugs, they must be used exactly the way they are prescribed
- Taking your treatment correctly is as important as which drugs you and your health care provider choose
- Before getting started, be prepared!
 - The right health care provider
 - Enough knowledge about HIV
 - A positive attitude



Starting Sooner Rather than Later

The results of a study called **the START trial** have shown once and for all:

 People living with HIV who start treatment early, while CD4 counts are still high, have a much lower risk of illness and death

Including people living with HIV who feel well

• Taking HIV drugs earlier reduced likelihood of developing AIDS-related *and* non-AIDS related illnesses



Consensus Statement

- Scientific experts and policy makers came together in July 2015
 - Issued statement declaring all people living with HIV should have access to HIV treatment as soon as they're diagnosed



Benefits of Starting Early

- A longer life
- Having a higher CD4 cell count and keeping it high
- Preventing further damage to the immune system
- Decreasing risk for HIV-related and non-HIV-related health problems
- Reducing risk of transmitting HIV to others (*Treatment* as Prevention or TasP)
- No risk of transmitting HIV to sexual partners if on treatment and undetectable viral load (*U=U*)



Risks of Starting Late

- A shorter life
- Severely weakened immune system
 - Can take longer to restore immune system to full strength
 - Delaying treatment can increase chances that people living with HIV will develop AIDS and other serious illnesses
- Greater chance of immune reconstitution syndrome
 - Can worsen infections when HIV drugs are started and CD4 count is low
- Passing HIV to others, including sexual partners and babies, if you become pregnant



HIV Treatment Guidelines

- Issued by a variety of global and national agencies
 - Written and reviewed regularly by HIV experts
- Guidelines on starting HIV treatment:
 - U.S. Department of Health and Human Services (DHHS): All people living with HIV should take HIV drugs, no matter their CD4 count
 - World Health Organization and others: Since September 2015 also recommend all people living with HIV should take HIV drugs, no matter their CD4 count



The Right Health Care Provider

- You and your provider ought to decide together when to start treatment
- Are you and your provider a team? Ask yourself:
 - Can I be totally honest with them?
 - Are they available when I have questions?
 - Do they take my concerns seriously?
- If so, great!



The Right Health Care Provider

- Not feeling fully respected? Consider talking to your provider or **finding another** (if you are able to choose)
 - Even if you have no choice of providers, being informed about
 HIV can help you in discussions with any provider
- Wise to have a health care provider who is an HIV specialist (treats many people living with HIV)
- To find HIV specialists near you:
 - Ask for suggestions from friends and/or ASOs
 - American Academy of HIV Medicine (AAHIVM)
 - HIV Medical Association (HIVMA)
 - International Association of Physicians in AIDS Care (IAPAC)



Goals of HIV Treatment

the well project

- Get your viral load as low as possible for as long as possible
- Preserve, improve health of immune system by increasing the number of CD4 cells
- Improve quality of life and reduce illness
- Reduce risk of transmitting HIV to others





Taking drugs on schedule, as prescribed

- To get best benefits from HIV therapy, good adherence is required
- HIV drugs need to be kept at a certain level in your body
- Skipping doses, not taking drugs on time, not following food requirements can cause drugs to be less effective or stop working



Resistance

When a drug can longer fight HIV effectively

- If viral load goes up while taking HIV drugs, you should have a resistance test
- If your HIV develops resistance, you will likely have to change some drugs in your regimen

The best way to avoid **resistance** is **to take your HIV drugs exactly as directed**!



A Positive Attitude

When you think about starting treatment, it's important to have a good attitude. May help to believe that:

- Starting treatment is the right decision for you
- HIV drugs will help you fight the virus
- You can take your medications the right way



A Positive Attitude

- Think about why you're starting treatment
 - To keep yourself in good health for your own sake, and/or for the good of your family
- You need not wait until your immune system is damaged before taking HIV drugs
- Newer HIV drugs are much easier on your body and have fewer side effects



A Positive Attitude

the **well** project

When you think about starting treatment, **it's important not to go it alone**

- Put together a **support system** that may include:
 - Your health care provider
 - Social workers
 - Case managers
 - Family and friends
 - Support group of other people living with HIV
 - Online communities such as The Well Project's A Girl Like Me!



Learn More!

- To learn more about this topic, please read the full fact sheet:
 - <u>Considerations Before Starting HIV Treatment</u>
- For more fact sheets and to connect to our community of women living with HIV, visit:
 - www.thewellproject.org
 - www.facebook.com/thewellproject
 - <u>www.twitter.com/thewellproject</u>