

thewellproject

HIV Treatment as Prevention (TasP)


Last updated: April 3, 2019

Together, we can change the course of the HIV epidemic...one woman at a time.

#onewomanatatime

www.thewellproject.org

#thewellproject




thewellproject

What Is Treatment as Prevention?

Treatment as prevention (TasP): using HIV treatment to **lower the risk of transmitting HIV**

- For people living with HIV:
 - Taking HIV drugs can reduce viral load, which means:
 - Less likely to transmit HIV to others
 - *Unable* to transmit HIV through sex (U=U)
 - Feel better because they have a healthier immune system






thewellproject

TasP to Prevent Perinatal Transmission


- In the 1990s, research showed pregnant women living with HIV who used Retrovir (**zidovudine** or **AZT**) greatly reduced risk of transmitting HIV to their babies
- Now standard practice to:
 - Test pregnant women for HIV
 - Provide HIV drugs to women with HIV during pregnancy, delivery, breastfeeding
 - Give HIV drugs to babies of women with HIV
- With HIV treatment and undetectable viral load, a woman's chance of transmitting HIV to her baby can be less than 1 in 100
- Called **preventing mother-to-child transmission** (PMTCT) in many parts of the world (term considered stigmatizing to many); also known as **preventing perinatal** or **vertical transmission**



thewellproject

TasP in Sexually Active Adults

- TasP can help **mixed-status couples** safely conceive
- **PrEP** can also help keep HIV-negative partner negative
- **2011 HTPN 052 study:** HIV treatment and undetectable viral load -> little chance of transmission to HIV-negative partner in mixed-status couple
 - Even when partner living with HIV had no HIV-related symptoms and CD4 count between 350-500 cells/mm³
- **2016 PARTNER study:** No transmission if partner living with HIV has undetectable viral load (**U=U**)




thewellproject

Treatment with a Public Health Twist

HPTN 052 *changed HIV prevention*

- Additional tool – treatment – in prevention toolbox
- How it works from **public health** (vs. individual health) point of view:
 - If enough people living with HIV could be treated so that their viral loads became undetectable
 - They would be much less likely to spread HIV
 - Number of new HIV cases would drop remarkably
 - Called a drop in **community viral load**




thewellproject

Treatment with a Public Health Twist

Test and treat strategies:


- Recommend universal HIV testing to identify, then treat, all people living with HIV (millions more than now get treatment)
 - Whatever their CD4 counts or viral loads
- Would require greatly expanded medical services
- Resource-limited countries may not be able to take on these additional public health tasks
 - Ethical concerns about who receives limited supplies of medical services and HIV drugs – those who need them for treatment, or those who need them for prevention, or both?



thewellproject

Treatment with a Public Health Twist

- ***How well TasP works can vary*** depending on factors including:
 - Health care systems' ability to provide testing, linkage to care, HIV drugs
 - Willingness, ability of people to get tested for HIV
 - HIV-related stigma, discrimination make people less likely to get tested
 - Willingness, ability of people living with HIV to adhere to HIV drug regimens so they remain virally suppressed




thewellproject

Other Methods for Preventing HIV

For HIV-negative people:

- **PrEP (Pre-Exposure Prophylaxis)**
 - Taking drugs before exposure to prevent disease
 - HIV-negative people taking HIV drugs to reduce their risk of getting HIV if exposed to the virus
 - Especially helpful for HIV-negative women who want to get pregnant and whose male partners are living with HIV
 - Promising tool women can use to ***prevent HIV without partners' knowledge or cooperation***




thewellproject

Other Methods for Preventing HIV

For HIV-negative people:

- **PEP (Post-Exposure Prophylaxis)**
 - Taking HIV drugs for +/- 1 month after possible HIV exposure (needle-stick, sex without condom, sexual assault)
 - To be effective, PEP must be taken right away – within first 72 hours after exposure to HIV, if possible




thewellproject

In Conclusion

Research continues to show ***TasP can be a game-changer*** in the course of HIV pandemic

TasP can not only positively affect **health of those living with HIV**, but also **protect those who are not living with HIV**
Undetectable = Untransmittable



thewellproject

Learn More!

- To learn more, and for links to articles featuring more details, please read the full fact sheet:
 - [HIV Treatment as Prevention \(TasP\)](#)
- For more fact sheets and to connect to our community of women living with HIV, visit:
 - www.thewellproject.org
 - www.facebook.com/thewellproject
 - www.twitter.com/thewellproject